

Red Giants Survey in ω Cen: preliminary FLAMES GTO results

Elena Pancino¹

INAF - Osservatorio Astronomico di Bologna
via Ranzani 1, I-40127 Bologna, Italy

Abstract. I present preliminary results for a sample of ~ 700 red giants in ω Cen, observed during the Ital-FLAMES Consortium GTO time in May 2003, for the Bologna Project on ω Cen. Preliminary Fe and Ca abundances confirm previous results: while the metal-poor and intermediate populations show a normal halo α -enhancement of $[\alpha/\text{Fe}] \simeq +0.3$, the most metal-rich stars show a significantly lower $[\alpha/\text{Fe}] \simeq +0.1$. If the metal-rich stars have evolved within the cluster in a process of self-enrichment, the only way to lower their α -enhancement would be SNe type Ia intervention.

1 Introduction

The treatment of the large data volumes that can be obtained with large telescopes and multi-objects spectrographs, such as FLAMES at the ESO VLT, requires a major upgrade of the methods commonly used to reduce and analyze high and medium resolution spectra.

We are therefore developing a set of routines for an automatic or semi-automatic abundance analysis of stellar spectra based on equivalent widths (EW). The first product is DAOSPEC, a code developed by P. B. Stetson for automatic EW measurement (<http://cadwww.hia.nrc.ca/stetson/daospec/>). The preliminary abundance analysis presented here is the first step of an interactive and automatic procedure under development at the Bologna Observatory.

2 Analysis and Results

Part of the Ital-FLAMES Guaranteed Time has been devoted to the study of ω Cen, within the framework of the Bologna Project on ω Cen [1]. 700 giants have been observed with the ESO-VLT GIRAFFE spectrograph in May 2003, with a resolution of $R \sim 20,000$ and a $S/N \simeq 50\text{--}150$ per pixel, spanning more than 4 magnitudes and covering all known sub-populations in the cluster. The spectral ranges observed (Setups HR09, HR13, HR11 and HR15) allow the determination of iron peak, α , s -process elements and of Eu.

The spectra have been reduced with the GIRAFFE BLDRS pipeline developed at the Geneva Observatory. EW have been measured with DAOSPEC [2], based on a linelist produced with the Vienna Atomic Line Database (VALD) [3]. Preliminary estimates of the stellar parameters T_{eff} , $\log g$, v_t and $[M/H]$ have been obtained from the WFI photometry published by [4] and the color-temperature calibration by [5]. MARCS model stellar atmospheres [6] have been




Fig. 1. $[\text{Ca}/\text{Fe}]$ versus $[\text{Fe}/\text{H}]$ for the ~ 700 stars observed with GIRAFFE. Typical errorbars are plotted in the lower-right corner. The typical disk-halo behaviour is represented by dotted lines.

employed together with the abundance analysis code by [7] to produce a first guess abundance of Fe and Ca, strictly based on the photometric parameter estimates.

The resulting $[\text{Ca}/\text{Fe}]$ versus $[\text{Fe}/\text{H}]$ plot is shown in Fig. 1, where except for a few outliers that will have to be manually inspected, a clear trend appears: $[\text{Ca}/\text{Fe}]$ slowly rises with $[\text{Fe}/\text{H}]$ until it reaches a maximum and then declines again for the most metal-rich stars (RGB-a according to [4]). This nicely confirms a previous finding by [8] and [9]. If the metal-rich stars have evolved within the cluster in a process of self-enrichment, the only way to lower their α -enhancement would be SNe type Ia intervention. No simple explanation is provided for the rise of $[\text{Ca}/\text{Fe}]$ at low $[\text{Fe}/\text{H}]$, although a series of star formation bursts should be the likely cause.

References

1. Ferraro, F. R., Bellazzini, M., Origlia, L., Pancino, E., & Sollima, A. 2003, ASP Conf. Ser. 296: New Horizons in Globular Cluster Astronomy, 215
2. Stetson, P. B. & Pancino, E., *in preparation*
3. Kupka, F., Piskunov, N., Ryabchikova, T. A., Stempels, H. C., & Weiss, W. W. 1999, A&AS, 138, 119
4. Pancino, E., Ferraro, F. R., Bellazzini, M., Piotto, G., & Zoccali, M. 2000, ApJ, 534, L83
5. Alonso, A., Arribas, S., & Martínez-Roger, C. 1999, A&AS, 140, 261
6. Edvardsson, B., Andersen, J., Gustafsson, B., Lambert, D. L., Nissen, P. E., & Tomkin, J. 1993, A&A, 275, 101
7. Spite, M. 1967, Annales d’Astrophysique, 30, 685
8. Pancino, E., Pasquini, L., Hill, V., Ferraro, F. R., & Bellazzini, M. 2002, ApJ, 568, L101
9. Origlia, L., Ferraro, F. R., Bellazzini, M., & Pancino, E. 2003, ApJ, 591, 916